


# El Niño and its Impacts on Federated States of Micronesia – Yap and Chuuk




### What is El Niño?

The El Niño – Southern Oscillation (ENSO) is a recurring climate pattern involving changes in the temperature of waters in the central and eastern tropical Pacific Ocean and the patterns of sea level pressure, lower- and upper-level winds, and tropical rainfall across the Pacific basin. On periods ranging from about two to seven years, the surface waters across a large swath of the tropical Pacific Ocean warm or cool by anywhere from 1°C to 3°C, compared to normal. This irregular oscillation between warm and cool patterns, referred to as the ENSO cycle, directly affects rainfall distribution in the tropics and can have a strong influence on weather across the Pacific basin. **El Niño** and **La Niña** are the extreme phases of the ENSO cycle; between these two phases is a third phase called **ENSO-neutral**.

**ENSO-neutral**: Under normal conditions strong trade winds blow from the east along the equator, pushing warm water into the western Pacific Ocean.



**El Niño** conditions occur when abnormally warm waters build in tropical region of the central and eastern Pacific Ocean and are usually associated with a weakening of the easterly trade winds, sometimes even reversing to westerlies. Consequently, tropical rains that usually fall over Indonesia move eastward; sea level decreases in the western Pacific; and the vertical, thermal structure of the ocean and coastal and upwelling currents are changed.



The *Thermocline* is a layer of water in which there is an abrupt change in temperature separating the warmer surface water from the colder deep water.

El Niño in FSM	
Rainfall	Less
more at first, but then very much less; longer and drier dry-season	•
Trade Winds	Less
weaker, with occasional westerly winds	•
Tropical Cyclones	Less
reduced risk, as more storms form closer to the Dateline	•
Sea Level	Less
lower at first, then gradually recovering	•
Ocean Conditions	Less
cooler at and below the surface	1
See back page for more details	

# **Every El Niño is a little bit different!**

**El Niño** conditions can start to develop as early as May or June and typically reaches maximum strength during December; the conditions then subside towards normal conditions by June of the following year. However, the evolution and duration, strength and impacts of individual El Niño events can vary, in some cases greatly. This makes constant monitoring and awareness extremely important for decision makers across multiple sectors.

# El Niño and Rainfall in Yap and Chuuk States

Rainfall during the El Niño year starts declining as soon as August in the western FSM. Dryness increases through October and November and peaks in January in the year after El Niño. Rainfall in western FSM can be as much as 50 to 60% below normal from January through April in the year after El Niño. The level of these dry conditions depends on the intensity of the El Niño event, with very strong El Niño events producing the driest conditions. Rainfall begins a return to normal by September of the year following an El Niño event, as long-term dryness lingers through June of the year after El Niño.

# **El Niño and Tropical Cyclones**

El Niño tends to increase the number of tropical cyclones (TC) eastward away from the Philippines and western Micronesia and towards the Dateline. This makes for increased risk of TC activity from March-July in the El Niño year, but a reduced risk from October-December.

### El Niño and Sea Level


During strong El Niño events, falls in sea level can be seen in Yap as early as February of the El Niño year and returning to normal as late as June of the year after El Niño. For moderate events, sea level usually starts falling during April or May and remains below normal until March or April of the following year.

## **El Niño and Ocean Conditions**

During El Niño, ocean waters tend to warm in the central and eastern Pacific at both the surface and subsurface. In FSM however, cooler than normal ocean waters are commonly observed during strong El Niño's, reducing tuna catch as many pelagic fish are displaced eastward.

# Yap Rainfall response to El Niño EL NINO YEAR YEAR AFTER EL NINO NORMAL N

Monthly rainfall during El Niño.



Monthly sea level variations during El Niño.

### What does El Niño mean to you?

- If you are a water manager, expect adverse impacts on water availability, including increased demand on groundwater resources as catchments and surface water sources become stagnant or dry up due to less rain.
- If you are a disaster manager, prepare for prolonged drought and a reduced likelihood of damage to infrastructure due to tropical cyclones and coastal flooding.
- If you are involved in public health, prepare for reduced freshwater quality and quantity, depleted food supplies, and mosquito-borne diseases.
- If you are a coastal and ocean resource manager, expect an increased risk of coral bleaching and potential changes in tuna catch as pelagic fish migrates away from its regular location.
- If you are involved in agriculture, expect adverse impacts on crop production and grass fires over the short term due to drought, and that these affects might linger as a result of seawater inundation of cropland.
- If you are involved with the recreation and tourism, drought and ecosystem impacts may negatively affect recreation and tourism.

### For Additional Information go to

- Weather Station Office (WSO) Yap or Chuuk: http://www.prh.noaa.gov/yap/ or http://www.prh.noaa.gov/chuuk/
- Pacific ENSO Applications Climate (PEAC) Center: http://weather.gov/peac/
- NOAA Climate Prediction Center (CPC): <a href="http://www.cpc.ncep.noaa.gov/">http://www.cpc.ncep.noaa.gov/</a>
- NOAA National Centers for Environmental Information (NCEI)

Also, Contact the Pacific Region Climate Officer,
Pacific ENSO Applications Climate Center, peac@noaa.gov







